

# We can prevent cervical cancer!

## There are simple steps to prevent cervical cancer.

### **Vaccinate early**

Cervical cancer vaccination is available for all genders and can be given as early as age nine. The vaccine gives the strongest immune response when given during the preteen years, but routine vaccination is recommended for everyone through age 26.

### **Screen regularly**

Cervical cancer screening should start no earlier than age 21. The specific approach to screening depends on the patient's age, medical history, and health care provider preference and may include a Pap test alone, an HPV test alone, or a Pap/HPV co-test. Ask your health care provider which approach is recommended for you and how often you should screen.

**Nearly all cases of cervical cancer are preventable!**



## Cervical Cancer Prevention in Pennsylvania

**496 new cases in  
Pennsylvania  
each year**

- Philadelphia: 85

**NCCC**

National Cervical Cancer Coalition  
A program of the American Sexual Health Association



## Your NCCC Philadelphia, Pennsylvania Chapter

Chapter Leader

Angela Reese

[areese@nccc-online.org](mailto:areese@nccc-online.org)

### Screening

The Affordable Care Act (ACA) requires insurance to cover cervical cancer screening.

Those lacking insurance can check with the Breast and Cervical Cancer Services program for help finding quality, low-cost breast and cervical cancer screening services.

You can find more information and eligibility requirements at:

### Pennsylvania Breast & Cervical Cancer Early Detection Program

Pennsylvania Department of Health  
Room 1011 Health and Welfare Building, Harrisburg, PA 17120  
Bucks, Chester, Delaware, Montgomery, and Philadelphia counties: call 800-848-3367 or 215-985-3300  
All other counties: call 800-215-7494

### Vaccination

Regardless of gender, there are a number of HPV-related diseases against which the vaccine protects. This is why HPV vaccines are available for everyone beginning at age nine.

**Why vaccinate so young?** The immune response to the vaccine is stronger in pre-teens, although it's still excellent in older youth and can be given routinely through age 26. Another advantage to early vaccination is only two doses are required until age 14 (those 15 and older need the full three-dose series).

## Cervical Cancer Screening Guidelines

Screening is not recommended for those under age 21 (cervical cancer is very rare in younger people). The specific approach chosen depends on factors such as age and medical history, and include:

- A Pap test alone every three years
- Co-testing with a Pap and HPV test, every five years
- An HPV test alone, every five years

Depending on the results of the Pap and/or HPV tests, a health care provider may recommend additional screening or procedures, so some people may be screened more often. Talk with your health care provider to see when screening should start (and how often), and which tests they recommend. Regardless of which approach is taken, the most important thing is to be screened regularly!



[www.nccc-online.org](http://www.nccc-online.org)